Photos needed

Hieraaetus pennatus (Dværgørn): 3rd year (open wing)

Ardea purpurea (Purpurhejre): 2nd year (front and back of an open wing)

Nycticorax nycticorax (Nathejre): 3rd year (front and back of an open wing)

Ardeola ralloides (Tophejre): 2nd year (front and back of an open wing)

Aquila chrysaetos (Kongeørn): 3rd year autumn/4th year spring (tail and front and back of an open wing); 5th year autumn/6th year spring (tail and front and back of an open wing)

Perdix perdix (Agerhøne): juvenile (open wing); post-juvenile (open wing)

Porzana porzana (Vandrikse): adult male in autumn after post-breeding moult (head, chin and open wing); adult female in autumn after post-breeding moult (head, chin and open wing)

Charadrius alexandrines (Hvidbrystet Præstekrave): female in spring (head); 2nd year in spring (open wing)

Charadrius morinellus (Pomeransfugl): 2nd year in spring (open wing) Calidris pugnax: 2nd year in spring (tail and open wing)

Calidris minuta (Dværgryle): 2nd year in spring (tail and open wing)

Calidris alpina (Alm. Ryle): male in spring (breast, upperparts and head); female in spring (breast, upperparts and head); 2nd year in spring (tail and open wing); adult in spring (tail and open wing).

Tringa erythropus (Sortklire): adult male in spring (crown and undertail coverts); adult female in spring (crown and undertail coverts); 2nd year in spring (tail and open wing)

Tringa nebularia (Hvidklire): 2nd year in spring (tail and open wing)

Glareola pratincole (Rødvinget Braksvale): juvenile (crown, upperparts, breast, tail and front and back of the open wing)

Strix aluco (Natugle): 4th year autumn (front and back of an open wing)

Lanius minor (Rosenbrystet Tornskade): 1st year in autumn (crown, head and open wing)

Monticola saxatilis (Stendrossel): male in autumn (upperparts, breast, tail an open wing); female in autumn (upperparts, breast, tail an open wing)

Turdus torquatus (Ringdrossel): juvenile (breast, upperparts, tail and open wing) Turdus pilaris: juvenile (breast, upperparts, tail and open wing) Saxicola rubetra: juvenile (breast, tail and open wing)

Emberiza hortulana (Hortulan): adult male in autumn (breast and open wing); adult female-like in autumn (breast and open wing

Authors mail contact is: jblascozum@gmail.com

Thank you very much,

Ariñe Crespo Diaz

aranzadi zientzia elkartea

SOCIEDAD DE CIENCIAS SCIENCE SOCIETY SOCIÉTÉ DE SCIENCES

